



**SUPPLEMENT No. 3**

TO

THE CYPRUS GAZETTE No. 2632 OF 4TH FEBRUARY, 1938.

**SUBSIDIARY LEGISLATION.**

---

**No. 13.**

**THE DANGEROUS DRUGS LAWS, 1925 TO 1932.**

**ORDER IN COUNCIL No. 1769**

**MADE UNDER SECTION 6 OF THE DANGEROUS DRUGS (AMENDMENT)  
LAW, 1932.**

**H. R. PALMER,**  
*Governor.*

Whereas it is enacted by section 6 of the Dangerous Drugs (Amendment) Law, 1932, that if His Excellency the Governor in Council thinks fit to declare that a finding with respect to any preparation containing any of the drugs, to which Part III of the Dangerous Drugs Laws, 1925 to 1932, applies, has, in pursuance of Article 8 of the International Opium Convention signed at Geneva on behalf of His Majesty on the nineteenth day of February, nineteen hundred and twenty-five, been communicated by the Council of the League of Nations to the Parties of the said Convention, the provisions of the said Part III shall as from such date as may be specified in the Declaration cease to apply to the preparations specified therein :

Now, therefore, His Excellency the Governor, with the advice of the Executive Council, is pleased to declare and it is hereby declared that findings with respect to the preparations specified in the Schedule hereto have in pursuance of Article 8 of the said Convention been communicated by the Council of the League of Nations to the parties to the said Convention and that the date from which the provisions of Part III of the Dangerous Drugs Laws, 1925 to 1932, shall cease to apply to the said preparations shall be the 13th day of June, 1937.

## SCHEDULE.

## (a) MORPHINE PREPARATIONS.

		In 1 bougie
1. <i>Cereoli iodoformi et morphinæ</i>	Iodoform .. .. .	0.320 gramme
	Morphine hydrochloride	0.016 „
	Oil of theobroma, sufficient to fill a 1-gramme mould.	
2. <i>Emplastrum opii.</i>	Elemi .. .. .	20 grammes
	<i>Terebinthina</i> .. .. .	30 „
	<i>Cera flava</i> .. .. .	15 „
	<i>Olibanum pulvis</i> .. .. .	18 „
	<i>Benzoes pulvis</i> .. .. .	10 „
	<i>Opii pulvis</i> .. .. .	5 „
3. <i>Emplastrum opii.</i>	<i>Balsamum peruvianum</i> ..	2 „
	Extract of opium .. .. .	25 grammes
	Refined elemi .. .. .	25 „
4. <i>Emplastrum opii.</i>	Diachylon plaster with gum .. .. .	50 „
	Elemi .. .. .	8 grammes
	<i>Terebinthina communis</i>	15 „
	<i>Cere flava</i> .. .. .	5 „
	<i>Olibani pulverata</i> .. .. .	8 „
	<i>Benzoes pulverata</i> .. .. .	4 „
	<i>Opii pulverati</i> .. .. .	2 „
	<i>Balsami peruviani</i> .. .. .	1 gramme
5. <i>Emplastrum opii.</i>	Opium, in very fine powder .. .. .	10 grammes
	Resin plaster .. .. .	90 „
	6. <i>Emplastrum opii</i> (see formula under 5) mixed with other plasters contained in the British Pharmacopœia or British Pharmaceutical Codex.	
7. <i>Linimentum opii.</i>	Tincture of opium .. .. .	500 millilitres
	Liniment of soap .. .. .	500 „
8. <i>Linimentum opii</i> (see formula under 7) mixed with any other liniment of the British Pharmacopœia or of the British Pharmaceutical Codex.		
9. <i>Linimentum opii ammoniatum.</i>	Ammoniated liniment of camphor .. .. .	30
	Tincture of opium .. .. .	30
	Liniment of belladonna	5
	Strong solution of ammonia .. .. .	5
	Liniment of soap to 100	
10. <i>Linimentum opii ammoniatum</i> (see formula under 9) mixed with any other British Pharmacopœia or British Pharmaceutical Codex liniment.		
11. <i>Caustic "Nerve Pastes."</i>	Preparations containing, in addition to morphine salts, or morphine and cocaine salts, at least 25 per cent. of arsenious acid, and made up with the requisite proportion of creosote or phenol to produce the consistency of a paste.	
12. <i>Diarrhœa Pills.</i>	Camphor .. .. .	0.0648 gramme
	Lead acetate .. .. .	0.013 „
	Bismuth subnitrate .. .. .	0.162 „
	Tannic acid .. .. .	0.0648 „
	Opium powder .. .. .	0.026 „

13. <i>Pilulæ digitalis et Opii compositæ.</i>	Digitalis leaves, in powder .. .. Opium in powder .. .. Ipecacuanha root, in powder .. .. Quinine sulphate .. .. Syrup of glucose, a sufficient quantity to make 12 pills.	0.31 gramme 0.19 „ 0.13 „ 0.78 „
14. <i>Pilulæ hydrargyri cum Opio.</i>	Mercury pill .. .. Opium, in powder .. .. To make 12 pills.	3.89 grammes 0.19 gramme
15. <i>Pilulæ hydrargyri cum Creta et Opii.</i>	Mercury with chalk .. .. Compound powder of ipecacuanha* .. .. Milk sugar, a sufficient quantity. Syrup of glucose, a sufficient quantity. To make 12 pills.	0.78 gramme 0.78 „
16. <i>Pilulæ ipecacuanhæ cum Scilla.</i>	Compound powder of ipecacuanha* .. .. Squill, in powder .. .. Ammoniacum, in powder .. .. Syrup of glucose, a sufficient quantity.	30 grammes 10 „ 10 „
17. <i>Pilulæ hydrargyri bichlorati cum Opii extracto.</i>	Bichloride of mercury triturated .. .. Extract of opium .. .. Extract of couch-grass .. .. Liquorice root in powder, <i>q.s.</i> for 10 pills.	10 centigrammes 20 „ 20 „
18. <i>Pilulæ hydrargyri iodati cum Opii pulvere.</i>	Hydrargyrum iodatum freshly prepared .. .. Opium powder .. .. Powdered liquorice .. .. White honey, <i>q.s.</i> for 10 pills.	50 centigrammes 20 „ 30 „
19. <i>Pilula plumbi, cum Opio.</i>	Lead acetate, in powder .. .. Opium, in powder .. .. Syrup of glucose .. .. (or a sufficient quantity)	80 grammes 12 „ 8 „
20. <i>Pilulæ terebinthine compositæ.</i>	Opium .. .. .. Chinini sulfas .. .. .. <i>Styrax liquidus</i> .. .. .. <i>Terebinthina laricina</i> .. .. .. <i>Magnesii subcarbonas</i> , a sufficient quantity to make 100 pills.	0.5 gramme 2 grammes 2 „ 8 „

\* The formula of this powder is given under 21, *Pulvis ipecacuanhæ compositus*.

21. <i>Pulvis ipecacuanhæ compositus</i> Syn.: <i>Pulvis ipecacuanhæ et opii</i> (Dover's powder).	Ipecacuanha root, in powder .. ..	10 grammes
	Opium, in powder .. ..	10 "
	Potassium sulphate in powder .. ..	80 "
22. Mixtures of <i>Dover's powder</i> (see formula under 21) with mercury and chalk, aspirin, phenacetin, quinine and its salts, and sodium bicarbonate.		
23. <i>Pulvis kino compositus</i> .	Kino, in powder .. ..	75 grammes
	Opium, in powder .. ..	5 "
	Cinnamon bark, in powder .. ..	20 "
24. <i>Suppositoria plumbi composita</i> . Syn.: <i>Suppositoria plumbi cum opio</i> .	Lead acetate, in powder .. ..	2.4 grammes
	Opium, in powder .. ..	0.8 gramme
	Oil of theobroma, a sufficient quantity for 12 suppositories, each weighing about 1 gramme.	
25. <i>Coryza Tablets No. 2</i> .	Powdered opium .. ..	0.0043 gramme
	Quinine sulph. .. ..	0.022 "
	Ammon. chlor. .. ..	0.022 "
	Camphor .. ..	0.022 "
	Ext. Belladonna leaves .. ..	0.0043 "
	Ext. aconite root .. ..	0.0043 "
26. <i>Diarrhœa Tablets No. 2</i> .	Powdered opium .. ..	0.016 gramme
	Camphor .. ..	0.016 "
	Powdered ipecacuanha .. ..	0.008 "
	Lead acetate .. ..	0.011 "
27. <i>Dysentery Tablets</i> .	Powdered opium .. ..	0.013 gramme
	Powdered ipecacuanha .. ..	0.0648 "
	Powdered calomel .. ..	0.0324 "
	Lead acetate .. ..	0.0324 "
	Bismuth betanaphthol .. ..	0.1944 "
28. <i>Tabella hydrargyri cum Opio</i> .	Mercurous chloride powder .. ..	0.065 gramme
	Antimony oxide powder .. ..	0.065 "
	Ipecacuanha-root powder .. ..	0.065 "
	Powdered opium .. ..	0.065 "
	Milk sugar .. ..	0.065 "
	Gelatine solution, a sufficient quantity to make 1 tablet.	
29. <i>Tabella plumbi cum Opio</i> .	Sugar of lead .. ..	0.195 gramme
	Powdered opium .. ..	0.065 "
	Gelatine solution, a sufficient quantity to make 1 tablet.	
30. <i>Tabletta plumbi cum Opio</i> .	Lead acetate, in fine powder .. ..	19.44 grammes
	Opium, in powder .. ..	3.24 "
	Refined sugar, in powder .. ..	6.48 "
	Ethereal solution of theobroma .. ..	3.60 mils
	Alcohol .. ..	0.90 mil

- |   |                           |    |
|---|---------------------------|----|
| 31. <i>Unguentum gallæ compositum</i> . | Galls in very fine powder | 20 |
|   | Extract of opium .. ..    | 4  |
|   | Distilled water .. ..     | 16 |
|   | Wool fat .. ..            | 10 |
|   | Soft paraffin, yellow ..  | 50 |
32. *Unguentum gallæ compositum* (see formula under 31) mixed with other ointments and plasters contained in the British Pharmacopœia or British Pharmaceutical Codex.
- |                                       |                       |              |
|---------------------------------------|-----------------------|--------------|
| 33. <i>Unguentum gallæ cum Opio</i> . | Gall ointment .. ..   | 92.5 grammes |
|                                       | Opium in powder .. .. | 7.5 "        |
34. *Unguentum gallæ cum Opio* (see formula under 33) mixed with other ointments and plasters contained in the British Pharmacopœia or British Pharmaceutical Codex.
35. *Yatren*—105 (Iodoxyquinoline-sulphonic acid) with 5 per cent. opium admixture.

## (b) COCAINE PREPARATIONS.

- |   |   |               |
|---|---|---------------|
| 1. <i>Bernatzik's Injections</i> .  | (a) <i>Hydrargyrum bicyanatum</i> .. .. .   | 0.03 gramme   |
|   | <i>Cocainum</i> .. .. .   | 0.02 "        |
|   | (b) <i>Hydrargyrum succinatum</i> .. .. .   | 0.03 "        |
|   | <i>Cocainum</i> .. .. .   | 0.01 "        |
| 2. <i>Stila's Injections</i> .  | (a) <i>Hydrargyrum succinatum</i> .. .. .   | 0.03 gramme   |
|   | <i>Cocainum muriaticum</i> .. .. .  | 0.01 "        |
|   | (b) <i>Hydrargyrum succinatum</i> .. .. .   | 0.05 "        |
|   | <i>Cocainum muriaticum</i> .. .. .  | 0.03 "        |
| 3. <i>Natrium bitoracicum compositum cum Cocaino</i> .  | In tablets, compressed tablets, lozenges, pastilles and the like, difficult to break up, and containing not more than 0.2 per cent. of cocaine salts in conjunction with not less than 20 per cent. borax and not less than 20 per cent. antipyrine, or some similar analgesic, and not more than 40 per cent. of flavouring matter. Maximum weight of each tablet, etc., 1 gramme. |               |
| 4. <i>Caustic "Nerve Pastes"</i> .  | Preparations containing in addition to cocaine salts or cocaine and morphine salts, at least 25 per cent. of arsenious acid, and made up with the requisite proportion of creosote or phenol to produce the consistency of a paste.   |               |
| 5. <i>Cocaine and Atropine Tablets</i> , with a content of not more than 0.0003 gramme of cocaine salts and not less than 0.0003 gramme of atropine salts to each tablet. | <i>Atropinum sulphuricum</i> .. .. .  | 0.0003 gramme |
|   | <i>Cocainum hydrochloricum</i> .. .. .  | 0.0003 "      |
|   | <i>Mannite</i> .. .. .  | 0.003 "       |
|   | Weight of one tablet .. .. .  | 0.0036 gramme |
|   | Cocaine content 8.3 per cent.   |               |

## (c) HEROIN PREPARATIONS.

1. <i>Elixir camphoræ compositum</i>	Camphor .. ..	4 grains
	Oil of anise .. ..	5 minims
	Benzoic acid .. ..	6 grains
	Diamorphine hydrochloride .. ..	4 "
	Liquid extract of ipecacuanha .. ..	120 minims
	Tincture of squill .. ..	1½ fl. ounce
	Simple syrup to 20 fl. ounces.	
2. <i>Elixir diamorphinæ et Terpini</i> , with <i>Apomorphine</i> .	Apomorphine hydrochloride .. ..	5 grains
	Diamorphine hydrochloride .. ..	4 "
	Terpin hydrate .. ..	44 "
	Alcohol .. ..	10 fl. ounces
	Glycerine .. ..	5 " "
	Syrup of wild cherry to 20 fl. ounces.	
3. <i>Linctus diamorphinæ</i> , with <i>Ipecacuanha</i> .	Liquid extract of ipecacuanha .. ..	120 minims
	Diamorphine hydrochloride .. ..	4 grains
	Tincture of hyoseyamus .. ..	1½ fl. ounce
	Spirit of chloroform .. ..	1½ " "
	Syrup of balsam of tolu .. ..	3 fl. ounces
	Syrup of wild cherry .. ..	3 " "
	Glycerine to 20 fl. ounces	
4. <i>Linctus senegæ compositus</i> .	Liquid extract of senega .. ..	1 fl. ounce
	Liquid extract of squill .. ..	1 " "
	Tartarated antimony .. ..	8 grains
	Diamorphine hydrochloride .. ..	4 "
	Glycerine .. ..	2 fl. ounces
	Simple syrup to 20 fl. ounces.	
5. <i>Linctus thymî compositus</i> .	Diamorphine hydrochloride .. ..	4 grains
	Apomorphine hydrochloride .. ..	5 "
	Distilled water .. ..	1 fl. ounce
	Liquid extract of thyme (I-I) .. ..	5 fl. ounces
	Solution of tolu .. ..	1¼ fl. ounce
	Glycerine to 20 fl. ounces.	

## (d) DICODIDE PREPARATIONS.

1. *Cardiazol-Dicodide Solutions*. Solutions containing not less than 10 per cent. of cardiazol and not more than 0.5 per cent. of dicodide salts.

## (e) EUCODAL PREPARATIONS.

1. <i>Anti-Opium Tablets.*</i>	Eucodal .. .. .	1 gramme
	<i>Pulvis gentianæ</i> .. .. .	35 grammes
	<i>Pulvis ipecacuanhæ</i> .. .. .	20 "
	Quinine sulphate .. .. .	20 "
	Caffeine .. .. .	5 "
	Sugar of milk .. .. .	25 "
	Mix up and make up 5-grain tablets.	
2. <i>Tablets B.B. Compound.</i>	<i>Berberis vulgaris</i> powder	0.0324 gramme
	<i>Nux vomica</i> .. .. .	0.013 "
	Eucodal .. .. .	0.0032 "
	Ipecacuanha .. .. .	0.0648 "
	Rhubarb .. .. .	0.013 "
	<i>Pulvis cinnamoni com-</i> <i>positus</i> .. .. .	0.0324 "
	Aromatic chalk .. .. .	0.0032 "

\* In exempting this preparation from the operation of the Geneva Convention, the Health Committee expressed the wish that it should not be offered to the public under the name of "anti-opium."

Ordered in Council, this 2nd day of February, 1938.

(M.P. 1456/20/2.)

R. J. THORNE THORNE,  
*Clerk of the Executive Council.*

## No. 14

## THE DANGEROUS DRUGS LAWS, 1925 TO 1932.

## ORDER IN COUNCIL No. 1770

## MADE UNDER SECTION 11 (2).

H. R. PALMER,  
*Governor.*

Whereas by section 11 (2) of the Dangerous Drugs Laws, 1925 to 1932, power is conferred on His Excellency the Governor by Order in Council to declare that Part III of the said Laws shall apply to any drug of whatever kind in the same manner as it applies to the drugs mentioned in sub-section (1) of the said section 11 if it appears to him that the drug is or is likely to be productive, if improperly used, or is capable of being converted into a substance which is or is likely to be productive, if improperly used, of ill effects substantially of the same character or nature as or analogous to those produced by Indian hemp, morphine or cocaine :

And whereas it appears to His Excellency the Governor that all preparations, admixtures or other substances (except syrupus Codeinæ Phosphatis B.P.C. 1934) containing any proportion of methylmorphine (commonly known as codeine) or ethylmorphine (commonly known as dionin) associated with an inert substance whether solid or liquid and all preparations, admixtures or other substances containing more than 2.5 per cent. of methylmorphine or ethylmorphine (calculated as pure